

5-11-2010

Comparative transcriptional profiling identifies takeout as a gene that regulates life span

Johannes Bauer

Southern Methodist University, jbauer@smu.edu

Michael Antosh

Brown University

Chengyi Chang

Brown University

Christoph Schorl

Brown University

Santharam Kolli

Southern Methodist University

See next page for additional authors

Follow this and additional works at: http://scholar.smu.edu/hum_sci_biosciences_research



Part of the [Biology Commons](#), and the [Genetics and Genomics Commons](#)

Recommended Citation

Bauer, Johannes; Antosh, Michael; Chang, Chengyi; Schorl, Christoph; Kolli, Santharam; Neretti, Nicola; and Helfand, Stephen L., "Comparative transcriptional profiling identifies takeout as a gene that regulates life span" (2010). *Biosciences Research*. 2.
http://scholar.smu.edu/hum_sci_biosciences_research/2

Authors

Johannes Bauer, Michael Antosh, Chengyi Chang, Christoph Schorl, Santharam Kolli, Nicola Neretti, and Stephen L. Helfand

Comparative transcriptional profiling identifies takeout as a gene that regulates life span

Johannes Bauer^{1,4,5}, Michael Antosh^{2,3,4}, Chengyi Chang¹, Christoph Schorl¹, Santharam Kolli⁵, Nicola Neretti^{1,2}, and Stephen L. Helfand¹

¹ *Department of Molecular Biology, Cell Biology and Biochemistry, Division of Biology and Medicine, Brown University, Providence, RI 02912, USA*

² *Institute for Brain and Neural Systems, Brown University, Providence, RI 02912, USA*

³ *Department of Physics, Brown University, Providence, RI 02912, USA*

⁴ *These authors shared equally in the work*

⁵ *Present address: Department of Biological Sciences, Southern Methodist University, 6501 Airline Drive, 237-DLS, Dallas, TX 75275, USA*

Running title: *Comparative transcriptional profiling identifies longevity gene*

Key words: *Dietary restriction, Calorie restriction, microarrays, Drosophila melanogaster, Sir2, p53, Rpd3, Indy, methuselah (mth), chico, life span extension, and takeout*

Correspondence: *Stephen L. Helfand, MD and Nicola Neretti. PhD, Department of Molecular Biology, Cell Biology and Biochemistry, Division of Biology and Medicine, Sidney Frank Life Sciences Building, Brown University, 185 Meeting Street, Providence, RI 02912, USA*

Received: 05/07/10; **accepted:** 05/09/10; **published on line:** 05/11/10

E-mail: Stephen_Helfand@Brown.edu; Nicola_Neretti@Brown.edu

Copyright: © Baue et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract: A major challenge in translating the positive effects of dietary restriction (DR) for the improvement of human health is the development of therapeutic mimics. One approach to finding DR mimics is based upon identification of the proximal effectors of DR life span extension. Whole genome profiling of DR in *Drosophila* shows a large number of changes in gene expression, making it difficult to establish which changes are involved in life span determination as opposed to other unrelated physiological changes. We used comparative whole genome expression profiling to discover genes whose change in expression is shared between DR and two molecular genetic life span extending interventions related to DR, increased dSir2 and decreased Dmp53 activity. We find twenty-one genes shared among the three related life span extending interventions. One of these genes, *takeout*, thought to be involved in circadian rhythms, feeding behavior and juvenile hormone binding is also increased in four other life span extending conditions: *Rpd3*, *Indy*, *chico* and *methuselah*. We demonstrate *takeout* is involved in longevity determination by specifically increasing adult *takeout* expression and extending life span. These studies demonstrate the power of comparative whole genome transcriptional profiling for identifying specific downstream elements of the DR life span extending pathway.

RESULTS

Genetic background affects the specific genes that respond to DR

We examined the relative change in gene expression under DR conditions in whole female flies at Days 10 and 40 using flies from a combined inbred yw/w^{1118} background and a Canton-S background. The DR conditions used (1.5N high calorie and 0.5N low calorie; [1]) extend life span by 30-40% in both of these backgrounds. Employing criteria of ≥ 1.5 fold change and ≤ 0.01 p value we found that the DR flies in the yw/w^{1118} background showed 1321 genes increased at Day 10 and 1140 genes decreased at Day 10 (Figure 1A). At Day 40 the yw/w^{1118} CR flies had only 129 genes increased and 19 genes decreased (Figure S1). In the Canton-S background 1286 genes increased with DR at Day 10 and 1435 genes decreased with DR at Day 10 (Figure 1A). At Day 40, 746 genes were increased and 715 genes were decreased in DR in the Canton-S background (Figure S1). Of the genes that in-

creased or decreased in DR at Day 10 approximately 55-60% (765 up; 708 down) of them were shared between the two different fly backgrounds (yw/w^{1118} and Canton-S). Gostat analysis of the genes altered by DR at day 10 and day 40 in these two different inbred genetic backgrounds revealed changes in biological functions similar to those previously described for DR in an outbred background of *Drosophila* [2-4]. (Table S1).

These studies indicate that by day 10 there are a substantial number of genes expressed differentially by DR flies: 2461 in yw/w^{1118} background; 2721 in Canton-S background and 1473 shared in both backgrounds. These changes should represent an inclusive set of most of the gene expression changes associated with DR including those unrelated to life span extension, but induced as a result of the nutritional challenge of DR. For example, in addition to extending life span in flies, DR also leads to a reduction in female fertility. The decrease in fertility is not thought to be a primary component of the life span extending effect [5-7].




Figure 1. Comparison of genes upregulated and downregulated in yw, w^{1118} DR, Canton-S CR and dSir2 overexpressed long-lived flies at Day 10. (A) Venn diagram comparing the upregulated and downregulated genes for DR flies in a yw/w^{1118} and a Canton-S background at age 10 days. (B) Venn diagram comparing upregulated and downregulated genes in DR long-lived flies and dSir2 overexpressing long-lived flies at age 10 Days. DR flies and dSir2 overexpressing flies are in an identical genetic background. The majority of genes expressed in dSir2 overexpression are also expressed in DR (72% upregulated and 61% downregulated). Verification of microarray data using qPCR is in Figure S3.

Gene expression changes in dSir2 overexpressing long-lived flies overlaps with DR long-lived flies

In order to identify genes involved more specifically in DR life span extension we compared the changes in gene expression in DR with those induced by a specific molecular genetic life span extending intervention related to DR that does not cause a decrease in female fertility; an increase in adult neuronal dSir2 expression [6]. To improve the sensitivity in detecting shared changes in gene expression in DR and dSir2 overexpressing flies we compared these two interventions in genetically identical flies by using the inducible RU486 system [8, 9]. A cohort of genetically identical flies possessing the GeneSwitch Elav driver (GSElav) and a construct permitting overexpression of dSir2 were randomly assigned to three different conditions: (i) high calorie food with EtOH diluent; (ii) low calorie food with EtOH diluent; and (iii) high calorie food with RU486.

A great deal of overlap in gene expression is seen between DR and neuronal specific dSir2 overexpression (Figure 1B). Of the 782 genes that change with neuronal specific dSir2 overexpression, 525 or 67% were shared with DR (72% upregulated and 61% downregulated). When the comparison is made between dSir2 overexpression and the genetically less related Canton-S DR the overlap is only 55% (428 genes out of 782—Figure S1).

Examination of the biological nature of the shared changes between DR and dSir2 life span extension at Day 10 using Gostat shows 148 shared categories decreasing and 72 shared categories increasing (a category contains at least 5 genes and a Gostat P value <0.05). The dSir2 long-lived flies share 78% of their downregulated and 72% of their upregulated GO categories with DR (Table S1). The comparison between DR and dSir2 overexpression also confirms the phenotypic observation that female reproduction is more significantly affected in DR than in dSir2 overexpression. DR downregulates 31 GO categories related to female reproduction, while dSir2 overexpression downregulates only 11 GO categories related to female reproduction (Table S1).

Gene expression changes in DN-Dmp53 expressing long-lived flies overlaps with DR and dSir2 overexpressing long-lived flies

Since a reduction in Dmp53 activity is a downstream component of the DR/Sir2 life span extending pathway [10, 11] we compared the changes in gene expression of DN-Dmp53 long-lived flies to DR and dSir2

overexpression in a similar genetic background. Examination of the changes in gene expression at Day 10 in flies expressing DN-Dmp53 revealed 132 genes are upregulated and 103 genes are down regulated (Figure 2A). Of the 235 genes that change with DN-Dmp53 expression, 87 or 37% were shared with DR (63% upregulated and 4% downregulated) and 88 or 37% were shared with dSir2 (65% upregulated and 2% downregulated) (Figure 2A). The relationship between changes in gene expression between DR, dSir2 and Dmp53 is illustrated by the heat map in Figure 2B. Only one shared gene is seen at Day 40 (Figure S1.)




Figure 2. Comparison of genes upregulated and down-regulated in DR, dSir2 overexpression and DN-Dmp53 expressing long-lived flies at Day 10. (A) Venn diagrams comparing upregulated and downregulated genes in DR, dSir2 overexpression, and DN-Dmp53 in a *yw/w¹¹¹⁸* background at age 10 Days. Genes intersecting in all 3 sets are noted in box with arrow. (B) Heatmap comparing the average log₂ fold changes for genes significantly altered in the *yw/w¹¹¹⁸* DR with the equivalent genes in dSir2 and DN-Dmp53 expressing flies.




Figure 3. *takeout* is overexpressed in several known life span extending conditions. *takeout* mRNA expression is increased in *yw/w¹¹¹⁸* DR, Canton-S DR, *w¹¹¹⁸* DR, *Sir2* overexpression, DN-Dmp53 expression, *Indy*, *Rpd3*, *methuselah* (*mth*) and *chico*. Fold change increase by qPCR of *takeout* mRNA from 10-Day old flies from these twelve life span extending conditions as compared to their genetically or dietary matched controls.

All but one of the 7 GO categories upregulated in the DN-Dmp53 expressing flies (endopeptidases, serine-type peptidase, serine-hydrolase, serine-type endopeptidase, peptidase, and defense response) are found in the upregulated GO categories of dSir2 and DR, while none of 15 GO categories downregulated in the DN-Dmp53 expressing flies are seen with dSir2 or DR (Table S1). Consistent with the normal fertility of the DN-Dmp53 expressing long-lived flies [5] we found no GO categories related to decreased female reproduction in the DN-Dmp53 expressing flies (Table S1).

Comparative whole genome expression profiling of DR, dSir2 and Dmp53 reveals a small set of shared genes

Comparison of the specific genes shared at Day 10 between these three related life span extending interventions (DR, dSir2 expression and DN-Dmp53 expression) show 20 genes upregulated and 1 gene down regulated (Figure 2A). Among the 20 upregulated genes are four genes associated with chromatin struc-

ture or maintenance (CG42249, CG5612, CG17325, CG4123), three genes associated with circadian rhythm (CG10553, CG13928 and *takeout*), two genes involved in neural activity (*Nplp3-neuropeptide-like precursor 3*, *synaptogyrin*), two genes involved in detoxification/chaperone activity (CG3091, CG6870), two genes involved in muscle maintenance (*Myo61F*, CG14687) and genes related to immune function (IM3-induced immune molecule 3), growth factor activity (*dawdle-activin*), and feeding behavior and response to starvation (*takeout*) [12]. The single downregulated gene is *Lsp2* (*larval serum protein-2*).

takeout is upregulated in other life span extending interventions

Of the 21 genes shared among the DR, dSir2 and DN-Dmp53 long-lived flies, *takeout* was the only gene significantly altered in transcriptional profiles of *Indy* long-lived flies [13]. We confirmed *takeout* was increased in *Indy* long-lived mutants by qPCR and found *takeout* to be increased in *Rpd3*, *chico*, and

methuselah mutants, single gene mutations that extend life span [14-16] (Figure 3). *takeout* was also found to be upregulated in DR in the Canton-S background and in an independent *w¹¹¹⁸* background by qPCR.

Increasing *takeout* expression extends life span

Given the association between the known phenotypes of *takeout* and longevity determination (feeding behavior,

response to starvation and juvenile hormone binding properties; [17-23]) and our finding of *takeout*'s upregulation in a number of different life span extending conditions, we examined the effect on life span of selectively increasing *takeout*. We found overexpression of *takeout* in adult neurons, pericerebral fat body or abdominal fat body extends male and female life span (Figure 4, Tables 1 and 2).




Figure 4. Overexpression of *takeout* in either of three different adult tissues extends life span of males and females. (A) Expression of *takeout* in the adult nervous system using the ELAV-Switch neuronal specific GAL4 driver leads to ~25% increase in mean longevity. (B) Flies expressing *takeout* in the head fat body, S1-32 pericerebral fat body specific GAL4 driver, have ~20% extension of mean life span, while *takeout* expression in the abdominal fat body, S1-106 abdominal fat body specific GAL4 driver, (C) extends fly life span by ~12-18% (females: left panel; males: right panel; statistical analysis in Table 1 and 2; grey: controls; black: *takeout*).

Table 1. The effect of *takeout* expression on female life span

| Driver | Mean LS (vs.ctrl) | Mean LS extension | Median LS (vs. ctrl) | Median LS extension | Max LS (vs. ctrl) | Max LS extension | Number of flies (control; experimental) | χ^2 | p-value |
|---------------------|-------------------|-------------------|----------------------|---------------------|-------------------|------------------|---|----------|---------|
| ELAV Switch | 48/44 | 9% | 48/44 | 9% | 64/60 | 7% | 275 248 | 12.92 | 0.0003 |
| ELAV Switch | 43/34 | 26% | 44/32 | 38% | 64/52 | 23% | 255 257 | 71.45 | <0.0001 |
| S ₁ -32 | 57/47 | 21% | 60/48 | 25% | 74/68 | 9% | 252 243 | 71.23 | <0.0001 |
| S ₁ -32 | 51/48 | 6% | 54/50 | 8% | 69/68 | 1% | 248 245 | 8.994 | 0.0027 |
| S ₁ -106 | 65/58 | 12% | 70/60 | 17% | 82/80 | 3% | 247 252 | 21.44 | <0.0001 |
| da | 50/40 | 25% | 50/40 | 25% | 66/56 | 18% | 256 247 | 119.3 | <0.0001 |

Log rank analysis of the survivorship curves of female flies. Mean, median and maximum lifespan, log rank analysis, p-value, percent change in mean, median and maximum lifespan as compared to controls (without RU486 for GeneSwitch experiments), Chi-square and p-values derived from the survivorship curves for each indicated intervention are shown. Maximum life span was calculated as the median life span of the longest surviving 10% of the population. ELAV Switch is the neuronal specific GeneSwitch GAL4 driver, S₁-32 is the pericerebral fat body specific GeneSwitch GAL4 driver, S₁-106 is the abdominal fat body specific GeneSwitch GAL4 driver and da is the *daughterless* GAL4 driver.

Table 2. The effect of *takeout* expression on male life span

| Driver | Mean LS (vs.ctrl) | Mean LS extension | Median LS (vs. ctrl) | Median LS extension | Max LS (vs. ctrl) | Max LS extension | Number of flies (control; experimental) | χ^2 | p-value |
|---------------------|-------------------|-------------------|----------------------|---------------------|-------------------|------------------|---|----------|---------|
| ELAV Switch | 53/52 | 2% | 56/54 | 4% | 66/64 | 3% | 228 190 | 1.34 | 0.247 |
| ELAV Switch | 50/43 | 16% | 52/42 | 24% | 70/62 | 13% | 241 228 | 31.08 | <0.0001 |
| S ₁ -32 | 50/46 | 9% | 54/46 | 17% | 66/60 | 10% | 234 234 | 18.34 | <0.0001 |
| S ₁ -32 | 66/63 | 5% | 66/64 | 3% | 84/84 | 0% | 247 243 | 1.669 | 0.1964 |
| S ₁ -106 | 52/44 | 18% | 54/48 | 13% | 64/60 | 7% | 227 233 | 81.9 | <0.0001 |
| da | 53/43 | 23% | 58/42 | 38% | 64/64 | 0% | 246 252 | 37.6 | <0.0001 |

Log rank analysis of the survivorship curves of male flies. Mean, median and maximum lifespan, log rank analysis, p-value, percent change in mean, median and maximum lifespan as compared to controls (without RU486 for GeneSwitch experiments), Chi-square and p-values derived from the survivorship curves for each indicated intervention are shown. Maximum life span was calculated as the median life span of the longest surviving 10% of the population. ELAV Switch is the neuronal specific GeneSwitch GAL4 driver, S₁-32 is the pericerebral fat body specific GeneSwitch GAL4 driver, S₁-106 is the abdominal fat body specific GeneSwitch GAL4 driver and da is the *daughterless* GAL4 driver.

Long-lived *takeout* overexpressing flies upregulate a subset of the genes upregulated in DR, dSir2 and DN-Dmp53 long-lived flies

It is our hypothesis that the twenty genes upregulated in flies whose life span is extended by DR, dSir2 or DN-Dmp53 may represent elements downstream in the DR life span extending pathway. Demonstration that upregulation of *takeout* results in life span extension confirms that *takeout* is a likely component of the DR life span extending pathway. As a first step in identifying additional downstream genes associated with DR life span extending pathways we examined which of the 19 remaining upregulated genes are also upregulated in long-lived *takeout* expressing flies using qPCR on- mRNA

from the flies overexpressing *takeout* in adult neurons. Nine out of the 19 genes showed a greater than 1.4 fold increase in expression in the *takeout* overexpressing long-lived flies (Figure 5). These include: (i) *dawdle*, a homologue of activin, coding for a transforming growth factor beta receptor binding protein; (ii) CG6870 coding for the cytochrome B5 detoxifying enzyme; (iii) CG3091, a gene coding for cellular retinaldehyde-binding/alpha-tocopherol transport that may be involved in detoxification; (iv) CG17325, a gene whose product interacts with chromatin related proteins such as SNR1; (v) CG42249, a gene with a predicted polycomb/trithorax response element; (vi) CG14687, with Myosin light chain binding properties; (vii) *Myo61F*; (ix) *synaptogyrin*; and (ix) CG5612 function unknown [12].




Figure 5. Long-lived *takeout* overexpressing flies have increased expression of a subset of the upregulated genes found in DR, dSir2 overexpression and DN-Dmp53 expressing flies. Overexpression of *takeout* in adult neurons increases the expression of a subset of 9 genes from the 19 upregulated genes shared in *yw/w¹¹¹⁸* DR, dSir2 overexpression and DN-Dmp53 expressing long-lived flies at Day 10. Fold change increase by qPCR of each of the noted genes using mRNA extracted from 10-Day old flies induced to express *takeout* (Elav-GeneSwitch-UAS-to) as compared to genetically identical controls Elav-GeneSwitch; UAS-to flies without fed diluent.

DISCUSSION

Examination of the changes in gene transcription profiles for DR in two different genetic backgrounds reveals the presence of a shared set of genes suggesting that one or more conserved core longevity-signaling pathways may exist to regulate lifespan in response to nutrient conditions. Such core longevity-signaling pathways may be utilized by other life span extending interventions not directly related to DR, and could help explain some of the cross-talk seen between DR and alterations in insulin/insulin-like signaling.

The set of common DR induced genes found represents genes important in life span extension as well as genes associated with other nutrient induced physiological functions not directly related to life span, such as decreased fertility. A comparative approach can be used to enrich for genes more specifically related to life span extension by examining life span extending interventions related to DR that do not have some of the same untoward effects as DR. Expression of dSir2 and DN-Dmp53 are two life span extending interventions that are part of the DR life span extending pathway in flies, but do not have decreased fertility [5, 6]. The whole genome expression profiles of flies on a DR diet and long-lived dSir2 expressing flies on a normal diet show a substantial overlap in changes in gene expression, supporting the observations linking dSir2 and DR (Figure 1B). As predicted, while DR has many GO categories associated with downregulation of fertility (31), fewer are seen with dSir2 long-lived flies (11) and none in DN-Dmp53 expressing long-lived flies (Table S1).

Comparisons of whole genome profiles of flies on DR, expressing dSir2 and expressing DN-Dmp53 revealed a small set of 21 commonly genes predicted to be enriched for genes involved in longevity regulation (Figure 2A). *takeout* (*to*), was selected to be further examined based upon *takeout*'s known role in regulating feeding behavior and the starvation response [17-23] as well as its presence in a set of upregulated genes from transcriptional profiles of another life span extending mutant in the fly, *Indy* [13]. Examination of *takeout* mRNA levels showed that in addition to *takeout* being upregulated in DR from three different fly backgrounds it is also upregulated in four additional separate life span extending mutants *chico*, *Rpd3*, *methuselah* and *Indy* [14-16, 24] (Figure 3). Confirmation of *takeout*'s role in longevity determination was demonstrated by overexpression in the fat body or nervous system of adult flies and extending life span (Figure 4).

The level of expression of *takeout* in the overexpression

studies is similar to the induction seen with DR (Figures 3 and S2), however lifespan extension by *takeout* overexpression is less than what is observed with DR. This effect may be due to the *w¹¹¹⁸* background used in these particular experiments, which is known to have a reduced DR response compared to other backgrounds [25]. Alternatively, *takeout* may be only one of several genes in the DR life span extending pathway that can positively influence lifespan. Other genes, including the additional 19 upregulated genes identified through comparative transcriptional profiling may increase lifespan incrementally, adding up to the lifespan extension total seen in DR or through other genetic interventions.

The mechanism by which increased *to* expression leads to life span extension is not known. Interestingly, *takeout* is regulated in a circadian fashion [18, 19, 26]. Increasingly, the link between the circadian system, food intake and aging has been observed [27]. The finding that expression of *takeout* from any of three different tissues (adult neurons, pericerebral fat body, abdominal fat body) extends life span suggests that the life span related functions of *takeout* could be due to its hypothesized function as a secreted Juvenile Hormone (JH) binding protein [17, 20, 22]. Although it is not known if the JH binding domain of *takeout* is functional, reduction of JH levels have been linked to increased longevity in grasshoppers [28]. *takeout* may bind JH in the hemolymph, thereby reducing JH bioavailability. It has been speculated that the insect ecdysone-JH system may be the functional equivalent of the mammalian thyroid hormone-prolactin axis, which controls important aspects of mammalian basal metabolism [29, 30]. Therefore, proteins such as *takeout* may be important mediators, linking a nutrient sensing network (DR, dSir2, insulin/insulin-like signaling) with an effector network (JH signaling), which in turn controls behavioral and physiological adaptation pathways.

Our data suggest that multi-factorial gene expression profiling can be successfully used to enrich for genes directly involved in the regulation of longevity, filtering out the noise of other physiological processes. Further refinement of this unbiased approach will be invaluable for discovering factors and signaling pathways involved in aging and lifespan regulation by a variety of modalities and for the identification of targets for specific therapeutic interventions.

EXPERIMENTAL PROCEDURES

All flies were kept in a humidified (50%), temperature-controlled incubator with 12 hour on/off light cycle at 25°C in vials containing standard cornmeal medium [6].

The ELAV-GeneSwitch line was from H. Keshishian (Yale University, New Haven, CT), S1-32-GeneSwitch, S1-106-GeneSwitch, *chico* and matched genetic controls for *chico* were from M. Tatar (Brown University, Providence, RI), *methuselah* and matched genetic controls from W. Ja (Caltech, Pasadena, CA) and UAS-*takeout* (UAS-*to*) was from B. Dauwalder (University of Houston, Houston, TX). All other lines (except *Indy*) were from the Bloomington *Drosophila* Stockcenter at Indiana University (Bloomington, IN).

The following crosses and experimental treatments were used in the microarray and lifespan analyses:

yw; ELAV-Geneswitch x P{EP}dSir2^{EP2300}/CyO (Bloomington 24859)

=> ELAV-Geneswitch-dSir2^{EP2300} (-/+ RU486) x P{GUS}-Dmp53^{259H}/TM6 (Bloomington 6582)

=> ELAV-Geneswitch-P{GUS}-Dmp53^{259H} (-/+ RU486)

yw; S1-32/CyO x UAS-*to*

=> S1-32-UAS-*to* (-/+ RU486)

yw; S1-106 x UAS-*to*

=> S1-106 -UAS-*to* (-/+ RU486)

Life span analysis. Flies were collected under light anesthesia, randomly divided into treatment groups and housed at a density of 25 males and 25 females each per vial. At least ten such vials were used per treatment as per [31]. Flies were passed every other day and the number of dead flies recorded.

All life span experiments were performed on regular cornmeal food, and for induction with the GeneSwitch system, RU486 (Sigma) was added directly to the food to a final concentration of 200µM. The same concentration of diluent (EtOH) was added to control food. RU486 was administered from the day of eclosion. For expression with constitutive da-GAL4 driver, UAS-*takeout* was backcrossed to w¹¹¹⁸ for 10 generations and isogenic controls were generated from the last backcross. Statistical analyses, including log rank tests, were performed using the Prism suit of biostatistical software (GraphPad, San Diego). Maximum life span was calculated as the median age of the last surviving 10% of the population.

Microarrays. For microarray experiments of DR animals, Canton-S and a mixed yw/w¹¹¹⁸ (the diluent controls from the genetic interventions below) line were aged for 10 or 40 days either on 1.5N or 0.5N food (15% sucrose and 15 yeast extract, or 5% sucrose and 5% yeast extract (all w/v), respectively) [1]. For genetic interventions, ELAV-GeneSwitch-dSir2^{EP2300} and

ELAV-GeneSwitch-DN-Dmp53^{259H} flies were aged for 10 or 40 Days as described for the life span experiments on food containing diluent or RU486. Total RNA was isolated from at least 75 females using Trizol (Invitrogen) and further purified using RNeasy columns (QIAGEN). 5 µg total RNA was used with Affymetrix One Cycle DNA conversion Kit (Cat # 900431) and all steps were carried out according to the Affymetrix manual. Briefly, first RNA was converted to double stranded cDNA followed by a clean-up step using spin columns. The double stranded cDNA was amplified in an in-vitro transcription reaction overnight at 37 °C using Affymetrix IVT labeling kit (cat # 900449), resulting in biotin labeled cRNA. After clean-up of the labeled cRNA with spin columns, 15 µg of cRNA were fragmented using metal induced hydrolysis. 10 µg of the fragmented RNA were hybridized to *Drosophila* 2.0 arrays overnight at 45 °C, 60 rpm. The array was stained using Affymetrix Hybridization-Wash-Stain kit and Fluidics Script FS450_0002 on the Affymetrix 450 fluidics station and finally, the arrays were scanned using an Affymetrix 3000 G7 scanner. At least three independent biological replicates per intervention were analyzed.

Pre-processing of microarray data: The data was quantile normalized and summarized using GCRMA [32] to obtain expression scores in the log2 scale. A probeset was considered absent if its mean expression level was below the 25th percentile (compared to the rest of the mean expressions for that condition) in both experiment and control. Absent probesets were removed from further analysis.

Differential Expression. A set of three biological replicates from both the treatment and control cohorts was used to identify differentially expressed probesets. Probesets with a p value (two sided t test) smaller than 0.01 and a fold change larger than 1.5 or smaller than 1/1.5 were selected as differentially expressed. These thresholds were chosen to minimize the number of false positives and false negatives in a comparison test of the microarray data of a pool of genes with PCR data from the same samples (Supplemental Figure 1). Probesets have been collapsed to genes after statistical selection for differential expression.

GOstat. The genes were analyzed using GOstat [4], which determines which sets of genes (called gene ontologies) are enriched in a list of genes. The input to GOstat is the list of differentially expressed genes for an experiment versus control comparison. For each gene ontology the intersection is found between the input list and the list of genes in the gene ontology. A p value is computed as the probability of obtaining an intersection

at least as large as the one observed by random sampling using the hypergeometric distribution. *p* values were adjusted for multiple testing using Benjamini and Hochberg's False Discovery Rate algorithm [33]. Ontologies with an adjusted *p* value < 0.05 were considered as overrepresented. Only gene ontologies containing at least 5 genes were considered.

Quantitative PCR. Total mRNA was isolated from at least 75 heads of 10-day old females using Trizol (Invitrogen) and further purified using the RNeasy kit (Qiagen). cDNA was generated with 0.5µg total mRNA in a 10µl reaction using the iScript cDNA synthesis kit (Bio-Rad). 0.8µl of the iScript reaction was used as qPCR template. qPCR was performed as described [10] on an ABI 7500 Real-Time PCR machine using the ABI SYBR-Green PCR master mix following the manufacturers instructions. Each qPCR reaction was performed using four biological replicates in triplicate each and normalized to mRNA from GAPDH or tubulin.

ACKNOWLEDGEMENTS

The authors would like to thank M. Tatar, W. Ja, B. Dauwalder, M. Roshbash and H. Keshishian for the kind gift of fly stocks and Will Lightfoot, Adam Kroll and Suzanne Hosier for technical assistance. This work was supported by NIA grants AG16667, AG24353 and AG25277 to SLH and NIA AG029723 to JHB, and NIA AG028753 to NN. MA was supported by NIA AG030329. The Brown Genomics Core is supported by NIH grants P20RR015578 and P20RR018728. SLH is an Ellison Medical Research Foundation Senior Investigator and recipient of a Glenn Award for Research in Biological Mechanisms of Aging.

CONFLICT OF INTERESTS STATEMENT

The authors of this manuscript have no conflict of interests to declare.


REFERENCES

1. Bross TG, Rogina B, and Helfand, SL. Behavioral, physical, and demographic changes in *Drosophila* populations through dietary restriction. *Aging Cell*. 2005; 4:309-317.
2. Pletcher SD, Libert, S, and Skorupa D. Flies and their golden apples: the effect of dietary restriction on *Drosophila* aging and age-dependent gene expression. *Ageing Res Rev*. 2005; 4:451-480.
3. Pletcher SD, Macdonald SJ, Marguerie R, Certa U, Stearns SC, Goldstein DB, and Partridge L. Genome-Wide Transcript Profiles in Aging and Calorically Restricted *Drosophila melanogaster*. *Curr. Biol*. 2002;12:712-723.
4. Beissbarth T and Speed TP. Gostat: find statistically overrepresented Gene Ontologies within a group of genes. *Bioinformatics*. 2004; 20:1464-1465.
5. Bauer JH, Poon PC, Glatt-Deeley H, Abrams JM, and Helfand SL. Neuronal expression of p53 dominant-negative proteins in adult *Drosophila melanogaster* extends life span. *Curr Biol*. 2005; 15:2063-2068.
6. Rogina B and Helfand SL. Sir2 mediates longevity in the fly through a pathway related to calorie restriction. *Proc Natl Acad Sci U S A*; 2004 101:15998-16003.
7. Grandison RC, Piper MD, and Partridge L. Amino-acid imbalance explains extension of lifespan by dietary restriction in *Drosophila*. *Nature*. 2009; 462:1061-1064.
8. Osterwalder T, Yoon KS, White BH, and Keshishian H. A conditional tissue-specific transgene expression system using inducible GAL4. *Proc. Natl. Acad. Sci. USA* ; 2001; 98: 12596-12601.
9. Roman G, Endo K, Zong L, and Davis RL. P[Switch], a system for spatial and temporal control of gene expression in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA*. 2001; 98:12602-12607.
10. Bauer JH, Morris SN, Chang C, Flatt T, Wood JG, and Helfand SL. dSir2 and Dmp53 interact to mediate aspects of CR-dependent life span extension in *D. melanogaster*. *Aging*. 2009; 1:38-48.
11. Donehower LA. Longevity regulation in flies: a role for p53. *Aging*. 2009; 1:6-8.
12. Tweedie S, Ashburner M, Falls K, Leyland P, McQuilton P, Marygold S, Millburn G, Osumi-Sutherland D, Schroeder A, Seal R, et al. FlyBase: enhancing *Drosophila* Gene Ontology annotations. *Nucleic Acids Res*. 2009; 37:D555-559.
13. Neretti N, Wang PY, Brodsky AS, Nguyen HH, White KP, Rogina B, and Helfand SL. Long-lived Indy induces reduced mitochondrial reactive oxygen species production and oxidative damage. *Proc Natl Acad Sci U S A*. 2009; 106:2277-2282.
14. Lin YJ, Seroude L and Benzer S. Extended life-span and stress resistance in the *Drosophila* mutant *methuselah*. *Science*. 1998; 282:943-946.
15. Rogina B, Helfand SL, and Frankel S. Longevity regulation by *Drosophila* Rpd3 deacetylase and caloric restriction. *Science*. 2002; 298:1745.
16. Clancy DJ, Gems D, Harshman LG, Oldham S, Stocker H, Hafen E, Leivers SJ, and Partridge L. Extension of life-span by loss of CHICO, a *Drosophila* insulin receptor substrate protein. *Science*. 2001; 292:104-106.
17. Meunier N, Belgacem YH, and Martin JR. Regulation of feeding behaviour and locomotor activity by takeout in *Drosophila*. *J Exp Biol*. 2007; 210:1424-1434.
18. Sarov-Blat L, So WV, Liu L, and Rosbash M. The *Drosophila* takeout gene is a novel molecular link between circadian rhythms and feeding behavior. *Cell*. 2000; 101:647-656.
19. So WV, Sarov-Blat L, Kotarski CK, McDonald MJ, Allada R, and Rosbash M. takeout, a novel *Drosophila* gene under circadian clock transcriptional regulation. *Mol Cell Biol*. 2000; 20:6935-6944.
20. Noriega FG, Ribeiro JM, Koener JF, Valenzuela JG, Hernandez-Martinez S, Pham VM, and Feyereisen R. Comparative genomics of insect juvenile hormone biosynthesis. *Insect Biochem Mol Biol*. 2006; 36:366-374.

21. Wong R, Piper MD, Wertheim B, and Partridge L. Quantification of food intake in *Drosophila*. *PLoS One*. 2009; 4:e6063.
22. Hamiaux C, Stanley D, Greenwood DR, Baker EN, and Newcomb RD. Crystal structure of Epiphyas postvittana takeout 1 with bound ubiquinone supports a role as ligand carriers for takeout proteins in insects. *J Biol Chem*. 2009; 284:3496-3503.
23. Lazareva, A.A., Roman, G., Mattox, W., Hardin, P.E., and Dauwalder, B. (2007). A role for the adult fat body in *Drosophila* male courtship behavior. *PLoS Genet*. 2007; 3:e16.
24. Rogina B, Reenan RA, Nilsen SP, and Helfand SL. Extended life-span conferred by cotransporter gene mutations in *Drosophila*. *Science*. 2000; 290:2137-2140.
25. Libert S, Zwiener J, Chu X, Vanvoorhies W, Roman G, and Pletcher SD. Regulation of *Drosophila* life span by olfaction and food-derived odors. *Science*. 2007; 315:1133-1137.
26. Benito J, Hoxha V, Lama C, Lazareva AA, Ferveur JF, Hardin PE, and Dauwalder B. The circadian output gene takeout is regulated by Pdp1epsilon. *Proc Natl Acad Sci U S A*. 2010; 107:2544-2549.
27. Imai SI. "Clocks" in the NAD World: NAD as a metabolic oscillator for the regulation of metabolism and aging. *Biochim Biophys Acta*. 2009; doi:10.1016/j.bbapap.2009.10.024
28. Tatar M and Yin C. Slow aging during insect reproductive diapause: why butterflies, grasshoppers and flies are like worms. *Exp Gerontol*. 2001; 36:723-738.
29. Davey K. From insect ovaries to sheep red blood cells: a tale of two hormones. *J Insect Physiol*. 2007; 53:1-10.
30. Marsh J. Developmental biology. Transforming frogs and flies. *Nature*. 1993; 361:116-117.
31. Bauer JH, Chang C, Morris SN, Hozier S, Andersen S, Waitzman JS, and Helfand SL. Expression of dominant-negative Dmp53 in the adult fly brain inhibits insulin signaling. *Proc Natl Acad Sci U S A*. 2007; 104:13355-13360.
32. Wu Z, Irizarry R, Gentleman R, Martinez Murillo F, and Spencer F. A model based background adjustment for oligonucleotide expression arrays. *Journal of the American Statistical Association*. 2004; 99:1-23.
33. Benjamini Y and Hochberg Y. Controlling the False Discovery Rate – A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. 1995; Series B (Methodological)* 57:289-300.



SUPPLEMENTAL DATA

The data of Supplemental Table 1 are found in full text version of this manuscript.



Supplemental Figure 1. Comparison of genes upregulated and downregulated in: *yw/w¹¹¹⁸* Dr, Canton-S background at age 40 Days. **(B)** Venn diagram comparing upregulated and down-regulated genes in DR long-lived flies and dSir2 overexpressing long-lived flies at age 40 Days. DR flies and dSir2 overexpressing flies are in an identical genetic background. Canton-S DR and dSir2 overexpressed long-lived flies at Day 10 and Day 40. Venn diagram comparing the upregulated and downregulated genes for DR flies in a Canton-S background and dSir2 overexpressing long-lived flies at age 10 Days **(C)** and age 40 Days **(D)**. Canton-S DR flies and dSir2 overexpressing flies are in different genetic backgrounds. DR, dSir2 overexpression and DN-Dmp53 expressing long-lived flies at Day 40. **(E)** Venn diagrams comparing upregulated and down-regulated genes in DR, dSir2 overexpression, and DN-Dmp53 in a *yw/w¹¹¹⁸* background at age 40 Days.

Supplemental Figure 2. *takeout* mRNA expression is increased in Elav GeneSwitch;UAS-to, S1-32; UAS-to and S1-106; UAS-to. Fold change increase by qPCR of *takeout* mRNA from 10-Day old flies from these three life span extending conditions as compared to their genetically matched controls.



Supplemental Figure 3. Verification of microarray data using qPCR. Each point on the graph represents a gene measured by both microarray and qPCR. The axes describe the fold change and p value of the microarray data. The red dots represent genes with a significant fold change (>20%) in PCR, and the blue dots represent genes with a non-significant fold change in PCR. The dotted lines define a box of the region where the PCR data is most likely to be significant--fold change>1.5 (0.58 in log2 space) and p value<0.01.